MISSOURI BOTANICAL GARDEN
PETER H. RAVEN LIBRARY

GEORGE ENGELMANN BOTANICAL NOTEBOOKS



Pagination Note:
Since many of the items lack a specific page number, the page number displayed online refers to the sequentially created number each item was given upon cataloging the materials.

Bonn June 1858

Mamillaria polyedra Milchsaft. die Milch besteht
aus einer Menge kleiner Kügelchen, die sich mit
Jod nicht färben, und einer kleinen Anzahl größerer
welche Öltröpfchen zu sein scheinen. Die ersten mögen
Harz sein, und die andern Cautschouk [illegible]. Kali (in Wasser
aufgelöst) hat keine Wirkung darauf. Auch wird sie durch
Ammoniak nicht, aber Äther löst sie rasch auf,
ebenso Alcohol

1110

Mamillaria Closiana Roum. in Bull. Soc. Bot. France II. 372

from the neighborhood of the City of Mexico. between M. Hystrix & M. Karwinskiana

1111

0 1 2 3 4 5 6 7 8 9 10
cm copyright reserved
MISSOURI BOTANICAL GARDEN

Angulosae a Tetragonae ——— crocidata. sempervivi

b. polyedrae xanthotricha polyedra
anisacantha
polygona

c. macrothelae ——— magnimamma
arietina. pentacantha

Phymatothelae Ludwigii macrothele. Lehmanni

Glanduliferae ——— cornifera,
Elephantidens.

Aulacothelae ———

Pelecyphora aselliformis is immersed!

Dr Kummer says that Mam. setosa, rhodanthe, & columnaris all belong to polythele.

1112

Munich Nov 1857

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI
BOTANICAL
GARDEN

0 1 2 3 4 5 6 7 8 9 10
cm copyright reserved
MISSOURI
BOTANICAL
GARDEN



1



2

white turned black!

MAMMILLARIA WRIGHTII.

Flower of Mam no 93

1113

H. B. Möllhausen

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI
BOTANICAL
GARDEN



Mamillaria flavescens, DC ?

Lima July 2d 1839

U.S. Exploring Exped.

Flowers subterminal?
is it a Coryphantha?

1114

0 1 2 3 4 5 6 7 8 9 10
cm copyright reserved
MISSOURI
BOTANICAL
GARDEN



1114

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI
BOTANICAL
GARDEN



18–22 white gracil
rad. spines
4 darker connivente
central ones.

proliferous branching
watery juice

M. discolor
M discolor Goebel
May
1860

Fl. 10–11 l. long with entirely exsert ovary! from a woolly axilla
20–22 linear-oblong petals, in several series,
obtuse, rose coloured in center
5 stigmata erect adpressed yellow
lower part of tube 1 line long
extremely narrow, naked.



1115

ALEX. LEITCH,
APOTHECARY & PHARMACEUTIST,
Marble Building Cor. Fourth & Olive Sts. St. Louis, Mo.

0 1 2 3 4 5 6 7 8 9 10
cm copyright reserved
MISSOURI
BOTANICAL
GARDEN

Mamillaria pusilla May 1862

pulp of fruit itself

x 30



scarcely any pulp in seminal cavity what there is seems to be composed entirely of the large cellular juicy funiculi – but seed very readily detached from them in ripe fruit.

Pits in the moist fresh seed not visible, appearing only when it gets dry – disappearing again upon moistening them.

1116

0 1 2 3 4 5 6 7 8 9 10 cm copyright reserved
MISSOURI BOTANICAL GARDEN

ALEX. LEITCH,
APOTHECARY & PHARMACEUTIST,
Marble Building Cor. Fourth & Olive Sts. St. Louis, Mo.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI
BOTANICAL
GARDEN

Mam Wildiana   June 1860

ovules 13/60 lines

tube elongated, contracted above naked ovary, naked at base
nearly 1 line





Pollen grains mostly
bluntly 3 angular 9/412 lin diameter
rarely 9½/412

1117

ALEX. LEITCH,
APOTHECARY & PHARMACEUTIST,
Marble Building, Cor. Fourth & Olive Sts., St. Louis, Mo.

Mam. Schiedeana May 1862

Cult St Louis



color best.

all the fruits sterile!

1118

M. W. ALEXANDER,
SOUTH-EAST CORNER OF FOURTH AND MARKET STREETS,
ST. LOUIS.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI
BOTANICAL
GARDEN

July 23 1861

Mamillaria Schelhaasii var rosea

from seeds recd at Mr Baumann's, Vienna, under the name of M. Scheidweileriana (no 105) Seedlings now in the third year, or 2¼ years old in flower since several weeks.

I notice some plants with smaller dirty whitish and some with larger red flowers; ~~all~~ from the same seed; the former are the regular M. Schelhaasii, the latter M Scheidweileriana or var rosea or purpurea of the former flowers of the former 7½ l. long, 6 lines wide, filaments very pale red — otherwise the same

M. Schelh. var rosea flowers 9 l. long. 8 lines in diameter when fully open.

Ovary exsert! oval,
tube constricted above the ovary, widening upwards; naked and green below.

Sepals 8, the inferior ones oblong-linear ciliate denticulate, mucronate, upper ones minutely serrate.

Petals 13 shorter than inner sepals and narrower, (acute linear) aristate

Lower part of tube constricted naked nectariferous, ¾ lines long

filaments red anthers almost orbicular-reniform.

Style longer - stigmata 3 oval ~~at last~~ patulous or at last reflexed.

Ovules rather few, on single very filamentose funiculi of their own length.

Mam. Schelhaasii var rosea
cult. ex [illegible]. Hort Baumann.
Vienna
sub nom M. Scheidweileriana
July 23 1861

x 412

$\frac{11\frac{1}{2}'''}{412}$



Pollen

x 30



x 30



older style

x 30



younger style

x 60



1120

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
Missouri
Botanical
Garden

M. W. ALEXANDER,
SOUTH-EAST CORNER FOURTH AND MARKET STREETS,
ST. LOUIS.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI
BOTANICAL
GARDEN

Mamillaria Heyderi var applanata
Texas

April 23 1861

1121


x 412
Pollen
x 30
x 60
x 30
x 4

seven greenish
yellow stigmata

copyright reserved
MISSOURI
BOTANICAL
GARDEN

M. W. ALEXANDER,
SOUTH-EAST CORNER FOURTH AND MARKET STREETS,
ST. LOUIS.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI
BOTANICAL
GARDEN

ex Hort Chalmers

parce *lactescens*

Mamill. coronaria var.

May 1861

6 dark central 16–18 white radial spines

flores ovario *emerso* in axilla parce villosa

5 stigm. yellow

9–10 sepala } all deep purple
15–18 petala mucronat lanceolat } all deep purple

x10



x4

x412

Pollen





abortive Pollen

x30



x60





x270

part of funiculus with the appendages

1122

M. W. ALEXANDER,
SOUTH-EAST CORNER FOURTH AND MARKET STREETS,
ST. LOUIS.

Bonn June 1858

Mamillaria polyedra schmutzig gelbe röthliche Blüthe 5–6 gelbe Stigmata, sepala länglich acuminat-aristat, ciliat, petala 8 kürzer, aristata, etwas lacerirt an der Spitze, — unterer Theil der Röhre, fast 1/2 Lin lang nackt

Pollen etwa 3 kantig  viel kleiner als bei Opuntia, sehr fein gekörnt; mit Schwefelsäure behandelt, welche an den drei Stellen eingeschnürt  oder ringsum geschlossen, sehr dünn.

Funiculus nicht immer haarig, mit sehr vielen und langen einzelligen Haaren, über die Micropyle gelagert, wie bei Cereus





filamenta gebräunt, antheren in verschiedener Stellung

Haare u. Wolle in der Axille ohne Intervall; — erstere langzellig spitz; innere äussere sepala fast.;

Mam. mutabilis

oblong linear stumpflich, beide sehr wenig und sehr fein gewimpert; petala 10–12 kürzer, spitz, Basis roth purpurn — stigmata gelblich weiß, 6 unterer Theil der Blüthe röhre nackt über ~~Mittelpunkt~~ 1/2 einer Linie lang.

1123

Ovula auf dem Rücken papillos, eigenthümlich funiculus die Micropyle bedeckend, [illegible] immer



Pollen ähnlich dem der vorigen
schwach rund in Wasser[illegible].
Die Haare an der Basis der Blüthe sind gleich denen der areola, glatt, gedreht nur sind die Zellen länger, wie das ganze Haar sehr lang ist.

Zellen auf der Oberfläche des Ovarium vieleckig, zum Theil leicht undeutlich. Stomata



Mam. Schiedeana Warzen mit horizontal abstehenden einzelligen Stachelhaaren oder Haaren bedeckt.

M. multiceps mit ähnlichen Warzen, doch sind sie oft länger haarig
Epidermis mit großen schlauchförmigen Zellen sowie Kern mit nucleolus  an feinen Fäden [illegible] dicht. — keine Doppelzellen

Aristolochia Sipho (und Althaea?) hat einen pubescirende Areola, da wo die Äste hervorkommen, in der axilla; diese Haare rundlich und 4-5 zellig, spitz.

Hort Schaefer, Münster June 1857

Mam. glochidiata var rosea, a very pretty cespitose plant, with red pulp, resembling in that respect M. pusilla; covered with pretty red flowers.

5 very small white stigmata, small naked space at the base of tube; 13 acute petals

1124

Mam. viridis with bright yellow flowers, but of same size than most similar Mamillariae

M. mutabilis and M. persicina have deep red crimson flowers.

M. quadrispina and M. recurva are a little lighter red.

M. persicina has a head of 5 green stigmata, very small naked space in base of tube of flower.

Mam. uberiformis in hort Poselg. Berlin May 30th 1857

ovarium exsertum; fl. 1 inch long and 1 inch diameter pale sulphur yellow, deep orangeyellow anthers 6-7 stigmata — interior sepals and outer petals pale, obtusish, denticulate, interior petals acuminate. — Very similar to M. sphaerica but flowers much smaller.

0 1 2 3 4 5 6 7 8 9 10 cm copyright reserved
MISSOURI BOTANICAL GARDEN